For Research Use Only

CSNK2A1 Monoclonal antibody

Catalog Number:68200-1-lg Featured Product

1 Publications



Basic Information

Applications

Catalog Number: 68200-1-lg Concentration:

1000 ug/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG29062

Tested Applications:

WB, IF/ICC, ELISA Cited Applications:

WB. IP

Species Specificity: human, rabbit Cited Species: human

GenBank Accession Number:

BC011668 GeneID (NCBI):

UNIPROT ID: P68400 Full Name:

> 45 kDa Observed MW: 40-45 kDa

Calculated MW:

casein kinase 2, alpha 1 polypeptide

Positive Controls:

WB: LNCaP cells, A549 cells, rabbit heart tissue, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, MOLT-4

Purification Method:

Protein G purification

Recommended Dilutions:

WB: 1:5000-1:50000 IF/ICC: 1:750-1:3000

CloneNo.:

1D5E8

cells, K-562 cells, HL-60 cells IF/ICC: A431 cells, HepG2 cells

Background Information

CSNK2A1(Casein kinase II subunit alpha) is also named as CK2A1 and belongs to the CK2 subfamily. The CSNK2A1 gene contains 8 exons whose sequences comprise bases 102 to 824 of the coding region of the human casein kinase II alpha subunit. Casein kinase II is a ubiquitous protein-serine/threonine kinase in eukaryotic cells that catalyzes the phosphorylation of many protein substrates and has been regarded as a second-messenger-independent protein kinase. This enzyme is composed of an α subunit (37 to 44kDa) and a β subunit (24 to 28 kDa) and exists as an ${\tt a}$ 2 ${\tt B}$ 2 tetramer (PMID:1448100). Casein kinase 2 has been found to be increased at protein level and up-regulated at the level of enzyme activity in the majority of cancers. However, elevated levels of casein kinase 2 are present in certain normal organs such as brain and testes.

Notable Publications

Author	Pubmed ID	Journal	Application
Riham Ayoubi	39403680	F1000Res	WB,IP


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68200-1-1g (CSNK2A1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



WB result of CSNK2A1 antibody (68200-1-1g; 1:10000; room temperature for 1.5 hours) with wild-type and sg-CSNK2A1 transfected A549 cells.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using CSNK2A1 antibody (68200-1-lg, Clone: 1D5E8) at dilution of 1:1500 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red).