

For Research Use Only

# EIF3B Monoclonal antibody

Catalog Number: 68202-1-Ig



## Basic Information

<b>Catalog Number:</b> 68202-1-Ig	<b>GenBank Accession Number:</b> BC001173	<b>Purification Method:</b> Protein G purification
<b>Size:</b> 1000 µg/ml	<b>GeneID (NCBI):</b> 8662	<b>CloneNo.:</b> 1G12G12
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> P55884	<b>Recommended Dilutions:</b> WB 1:2000-1:10000
<b>Isotype:</b> IgG1	<b>Full Name:</b> eukaryotic translation initiation factor 3, subunit B	
<b>Immunogen Catalog Number:</b> AG31427	<b>Calculated MW:</b> 93 kDa	
	<b>Observed MW:</b> 115 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : HCT 116 cells, A431 cells, HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells
<b>Species Specificity:</b> Human, mouse, rat	

## Background Information

EIF3B, also named Eukaryotic translation initiation factor 3 subunit B, is an 814 amino acid protein, which contains 1 RRM (RNA recognition motif) domain and 8 WD repeats and belongs to the eIF-3 subunit B family. EIF3B as an RNA-binding component of the eukaryotic translation initiation factor 3 (eIF-3) complex, is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation, and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression. The calculated molecular weight of EIF3B is 93 kDa, but the phosphorylated EIF3B protein is about 115 kDa.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68202-1-Ig (EIF3B antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.