

# KIAA2013 Monoclonal antibody

Catalog Number: 68245-1-Ig

## Basic Information

<b>Catalog Number:</b> 68245-1-Ig	<b>GenBank Accession Number:</b> BC035033	<b>Purification Method:</b> Protein A purification
<b>Concentration:</b> 1000 ug/ml	<b>GeneID (NCBI):</b> 90231	<b>CloneNo.:</b> 4G6C4
<b>Source:</b> Mouse	<b>UNIPROT ID:</b> Q8IYS2	<b>Recommended Dilutions:</b> WB 1:5000-1:50000 IF/ICC 1:500-1:2000
<b>Isotype:</b> IgG2b	<b>Full Name:</b> KIAA2013	
<b>Immunogen Catalog Number:</b> AG27438	<b>Observed MW:</b> 65 kDa	

## Applications

<b>Tested Applications:</b> WB, IF/ICC, ELISA	<b>Positive Controls:</b>
<b>Species Specificity:</b> human, rat	<b>WB :</b> A549 cells, LNCaP cells, HeLa cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells
	<b>IF/ICC :</b> HepG2 cells,


## Background Information

KIAA2013, also known as Q8IYS2 or MGC33867, is a single-pass transmembrane protein encoded by the KIAA2013 gene in humans. KIAA2013 has been found to play a role in the Endocannabinoid system. This system is made up of cannabinoid receptors 1 and 2 (CB1 and CB2) as well as the various ligands and enzymes that interact. The protein KIAA2013 has been found to be expressed within CB2 expressing cells. Both cannabinoid receptors are labeled as class A G-Protein Coupled Receptors, and CB2 is highly expressed within the human spleen and leukocytes. CB2, and by extension KIAA2013, are therefore targets of interest for therapeutic studies looking into diseases such as inflammatory bowel disease and rheumatoid arthritis(PMID:31616248 34752341).


## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
**Aliquoting is unnecessary for -20°C storage**


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using KIAA2013 antibody (68245-1-Ig, Clone: 4G6C4) at dilution of 1:1000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).



Various lysates were subjected to SDS PAGE followed by western blot with 68245-1-Ig (KIAA2013 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68245-1-Ig (KIAA2013 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.