

ABL1 Monoclonal antibody

Catalog Number: 68254-1-Ig

Basic Information

Catalog Number:

68254-1-Ig

Size:

1000 ug/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG26152

GenBank Accession Number:

NM_005157

GeneID (NCBI):

25

Full Name:

c-abl oncogene 1, receptor tyrosine kinase

Calculated MW:

123 kDa

Observed MW:

130 kDa

Purification Method:

Protein A purification

CloneNo.:

1E7A5

Recommended Dilutions:

WB 1:2000-1:10000

IF/ICC 1:400-1:1600

Applications

Tested Applications:

WB, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

WB : LNCaP cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells

IF/ICC : U2OS cells, HepG2 cells, MCF-7 cells

Background Information

ABL1, also named as ABL, JTK7, c-ABL and p150, belongs to the protein kinase superfamily, Tyr protein kinase family and ABL subfamily. ABL1 regulates cytoskeleton remodeling during cell differentiation, cell division and cell adhesion. ABL1 localizes to dynamic actin structures, and phosphorylates CRK and CRKL, DOK1, and other proteins controlling cytoskeleton dynamics. It regulates DNA repair potentially by activating the proapoptotic pathway when the DNA damage is too severe to be repaired. ABL1 catalyze the reaction: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate. A chromosomal aberration involving ABL1 is a cause of chronic myeloid leukemia (CML) [MIM:608232]. Translocation t(9;22)(q34;q11) with BCR. The translocation produces a BCR-ABL found also in acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL). The antibody is specific to ABL1.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68254-1-Ig (ABL1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Methanol) fixed HepG2 cells using ABL1 antibody (68254-1-Ig, Clone: 1E7A5) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using ABL1 antibody (68254-1-Ig, Clone: 1E7A5) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using ABL1 antibody (68254-1-Ig, Clone: 1E7A5) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10⁶ K-562 cells were intracellularly stained with 0.4 ug Anti-Human ABL1 (68254-1-Ig, Clone:1E7A5) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).