For Research Use Only

SCN5A Monoclonal antibody, PBS Only



Purification Method:

Protein G purification

CloneNo.:

1C2B3

Catalog Number: 68273-1-PBS

Basic Information

Catalog Number:

68273-1-PBS

Size:
1mg/ml
Source:
Mouse

Isotype:

Immunogen Catalog Number:

immunogen Catalog Number:

AG19275

19275

Observed MW: 227 kDa

BC140813

6331

Q14524

Full Name:

V. alpha subunit

Calculated MW: 2016 aa, 227 kDa

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

sodium channel, voltage-gated, type

Tested Applications: Indirect ELISA, IHC, WB

Species Specificity: rabbit, rat, mouse, human

Applications

Background Information

Voltage-gated sodium channels are responsible for initiation and propagation of action potentials in the membranes of neurons and most electrically excitable cells (PMID: 10798388). These channels are composed of a large alpha subunit that forms the ion conduction pore and auxiliary beta subunits (PMID: 11486343). The alpha subunits form a gene family with at least 10 members. Nav1.5, encoded by the SCN5A gene in humans, is a pore forming alpha subunit of voltage-gated sodium channels. Nav1.5 is the principal Na+ channel isoform expressed in cardiomyocytes. Mutations in SCN5A gene have been linked to many cardiac electrical disorders, including the congenital and acquired long QT syndrome, Brugada syndrome, conduction slowing, sick sinus syndrome, atrial fibrillation, and dilated cardiomyopathy (PMID: 23123192).

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


Selected Validation Data




rabbit heart tissue were subjected to SDS PAGE followed by western blot with 68273-1-lg (SCN5A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68273-1-PBS in a different storage buffer formulation.



rat heart tissue were subjected to SDS PAGE followed by western blot with 68273-1-lg (SCN5A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68273-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 68273-1-Ig (SCN5A antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68273-1-PBS in a different storage buffer formulation.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human SCN5A (68273-1-lg, Clone:1C2B3) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68273-1-PBS in a