For Research Use Only

DHX32 Monoclonal antibody

Catalog Number:68306-1-lg Featured Product

1 Publications

BC002473

55760

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:



Basic Information

Catalog Number: 68306-1-lg Concentration: 1000 μg/ml

Mouse Q7L7V1 Full Name: Isotype: lgG2b DEAH (Asp-Glu-Ala-His) box

Immunogen Catalog Number:

AG13956

Source:

polypeptide 32 Calculated MW: 743 aa, 84 kDa

Observed MW: 84 kDa

Purification Method: Protein A purification

CloneNo.: 2A4G9

Recommended Dilutions:

WB: 1:2000-1:10000

FC (Intra): 0.80 ug per 10^6 cells in a

100 µl suspension

Applications

Tested Applications: WB, ELISA, FC (Intra) **Cited Applications:**

Species Specificity: Human, Mouse, Rat **Cited Species:** human

Positive Controls:

WB: A431 cells, U2OS cells, rat testis tissue, HeLa cells, A549 cells, mouse testis tissue, MCF-7 cells, K-

562 cells

FC (Intra): HeLa cells,

Background Information

DHX32, an RNA helicase, belongs to the DEAH box RNA helicase family by virtue of its homology to the conserved $he licase\ domain.\ DHX32\ is\ abundantly\ expressed\ in\ human\ and\ murine\ models,\ participating\ in\ cell\ transcription$ and differentiation, and being capable of regulating T-cell apoptosis. DHX32 is also associated with some malignant tumors, such as lymphoma, colorectal cancer, and breast cancer (PMID: 35117530). DHX32 has 2 isoforms with molecular weights of 75 and 84 kDa, respectively.

Notable Publications

Author Pubmed ID Journal Application Mai Morita 40128189 Nat Commun

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68306-1-1g (DHX32 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of DHX32 antibody (68306-1-lg; 1:6000; incubated at room temperature for 1.5 hours) with sh-Control and sh-DHX32 transfected A549 cells.



1x10^6 HeLa cells were intracellularly stained with 0.8 ug DHX32 Monoclonal antibody (68306-1-Ig, Clone:2A4G9) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.8 ug Mouse IgG2b isotype control Mouse McAb (66360-3-Ig, Clone: 11B8C4) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).