For Research Use Only

HDDC2 Monoclonal antibody, PBS Only **proteintech***

Catalog Number:68391-1-PBS


Featured Product




Basic Information	Catalog Number: 68391-1-PBS	GenBank Accession Number: BC003357	Purification Method: Protein A purification
	Concentration: 1mg/ml	GenelD (NCBI): 51020	CloneNo.: 2E1B2
	Source: Mouse	UNIPROT ID: Q7Z4H3	
	Isotype: IgG2a	Full Name: HD domain containing 2	
	Immunogen Catalog Number: AG33258	Calculated MW: 23 kDa	
		Observed MW: 23 kDa	
Applications	Tested Applications: WB, IHC, IF/ICC, FC (Intra), Indirect ELISA		
	Species Specificity: human, mouse, rat		
Background Information	HDDC2, also named as C6orf74, plays a role in the maintenance of pluripotency in human ES and iPS cells. HDDC2 is a nuclear - encoded but mitochondrially localized protein. It has been reported as associated with oxidative metabolism.		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	yat-80°C

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 68391-1-lg (HDDC 2 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68391-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 68391-1-1g (HDDC2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68391-1-PBS in a different storage buffer formulation.



WB result of HDDC2 antibody (68391-1-Ig; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HDDC2 transfected HeLa cells. This data was developed using the same antibody clone with 68391-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Methanol) fixed Saos-2 cells using HDDC2 antibody (68391-1-Ig, Clone: 2E1B2) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68391-1-PBS in a different storage buffer formulation.



1x10^6 U2OS cells were intracellularly stained with 0.4 ug HDDC2 Monoclonal antibody (68391-1-lg, Clone:2E1B2) and Coralite@488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.4 ug Mouse IgG2a isotype control Mouse MCAb (66360-2-Ig, Clone: 11A1B2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68391-1-PBS in a different storage buffer