

For Research Use Only

GATC Monoclonal antibody, PBS Only

Catalog Number: 68405-1-PBS



Basic Information

Catalog Number: 68405-1-PBS	GenBank Accession Number: BC107145	Purification Method: Protein A purification
Size: 1mg/ml	GeneID (NCBI): 283459	CloneNo.: 2F9C8
Source: Mouse	UNIPROT ID: O43716	
Isotype: IgG2b	Full Name: glutamyl-tRNA(Gln) amidotransferase, subunit C homolog (bacterial)	
Immunogen Catalog Number: AG19857	Calculated MW: 136 aa, 15 kDa	
	Observed MW: 15 kDa	

Applications

Tested Applications:
WB, IF/ICC, Indirect ELISA

Species Specificity:
human

Background Information

GATC encodes a subunit of the glutamyl-tRNA(Gln) amidotransferase protein complex (GatCAB), which plays a role in aminoacylation of mitochondrial glutamyl mt-tRNA. The complex plays a role in mitochondrial translation and protein synthesis (PubMed: 30283131).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




1x10⁶ HEK-293 cells were intracellularly stained with 0.25 ug GATC Monoclonal antibody (68405-1-Ig, Clone:2F9C8) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1)(red), or 0.25 ug Mouse IgG2b isotype control Mouse McAb (66360-3-Ig, Clone: 11B8C4) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 68405-1-PBS in a different storage buffer



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using GATC antibody (68405-1-Ig, Clone: 2F9C8) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 68405-1-PBS in a different storage buffer formulation.



A549 cells were subjected to SDS PAGE followed by western blot with 68405-1-Ig (GATC antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68405-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 68405-1-Ig (GATC antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68405-1-PBS in a different storage buffer formulation.