### For Research Use Only

# **ULK1** Monoclonal antibody

Catalog Number:68445-1-lg 2 Publications



**Basic Information** 

Catalog Number:

GenBank Accession Number:

Purification Method:

68445-1-lg

NM\_003565

Protein G purification

Size:

GeneID (NCBI):

CloneNo.:

1000 µg/ml

8408

1D7G1

Source: Mouse UNIPROT ID: 075385 Recommended Dilutions:

Isotype:

Full Name:

WB 1:5000-1:50000 IHC 1:50-1:500

lgG1

unc-51-like kinase 1 (C. elegans)

Immunogen Catalog Number: AG22397

Observed MW: 130-150 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, ELISA
Cited Applications:

WB

WB: A549 cells, PC-3 cells, HeLa cells, Jurkat cells, HT-1376 cells, LnCaP cells, T-47D cells, NIH/3T3 cells, PC-

12 cells

Species Specificity:

Human, Mouse, Rat

Cited Species:

human, mouse

IHC: human liver tissue, mouse skeletal muscle tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

## **Background Information**

ULK1, also named as KIAA0722, belongs to the protein kinase superfamily, Ser/Thr protein kinase family and APG1/unc-51/ULK1 subfamily. It is involved in autophagy. ULK1 is required for autophagosome formation. It forms a stable complex with Atg13 and focal adhesion kinase (FAK) family interacting protein of 200 kDa (FIP 200) (PMID:21460634). ULK1 phosphorylates ATG13/KIAA0652. It is involved in axon growth. ULK1 plays an essential role in neurite extension of cerebellar granule cells. ULK1 represents a potential novel prognostic biomarker for HCC patients and may play an important role during the progression of HCC(PMID:23573318).

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Rui Qi Wang	39437445	Am J Physiol Cell Physiol	WB
Xiao Jin	39047963	ACS Infect Dis	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 68445-1-1g (ULK1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 68445-1-Ig (ULK1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 68445-1-Ig (ULK1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 68445-1-Ig (ULK1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 68445-1-Ig (ULK1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).