For Research Use Only

UNC13A Monoclonal antibody, PBS Only



Catalog Number: 68483-1-PBS

Basic Information

Catalog Number:

68483-1-PBS

Size: 1 mg/ml Source:

Mouse Isotype: IgG2a GenBank Accession Number:

NM_001080421

GeneID (NCBI):

23025 UNIPROT ID:

Q9UPW8
Full Name:

unc-13 homolog A (C. elegans)

Calculated MW: 193 kDa Observed MW: 193-210 kDa Purification Method:

Protein A purification CloneNo.:

3C3A9

Applications

Tested Applications: WB,IHC,IF,ELISA Species Specificity:

Human, mouse, rat, pig, rabbit

Background Information

UNC13A, also named KIAA1032, belongs to the unc-13 family. It plays a role in vesicle maturation during exocytosis as a target of the diacylglycerol second messenger pathway. UNC13A is involved in neurotransmitter release by acting in synaptic vesicle priming prior to vesicle fusion and participates in the activity-dependent refilling of readily releasable vesicle pool (RRP). It is essential for synaptic vesicle maturation in most excitatory/glutamatergic but not inhibitory/GABA-mediated synapses. UNC13A is also involved in secretory granule priming in INS secretion. The antibody is specific to UNC13A.

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 68483-1-lg (UNC13A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68483-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 68483-1-Ig (UNC 13A antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68483-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 68483-1-Ig (UNC 13A antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 68483-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed SH-SY5Y cells using UNC 13A antibody (68483-1-lg, Clone: 3C3A9) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 68483-1-PBS in a different storage buffer formulation.