

For Research Use Only

OTUB1 Monoclonal antibody

Catalog Number: 68489-1-Ig



Basic Information

Catalog Number: 68489-1-Ig	GenBank Accession Number: BC007519	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 55611	CloneNo.: 2F7F8
Source: Mouse	UNIPROT ID: Q96FW1	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2a	Full Name: OTU domain, ubiquitin aldehyde binding 1	
Immunogen Catalog Number: AG33132	Calculated MW: 31 kDa and 35 kDa	
	Observed MW: 31-35 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : U2OS cells, LNCaP cells, HeLa cells, HEK-293 cells, K-562 cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells
Species Specificity: Human, Mouse, Rat	

Background Information

OTUB1 belongs to the ovarian tumor domain protease (OTU) subfamily of deubiquitinases, which could block ubiquitination leading to protein stability. OTUB1 mainly prefers k-48 and k-63 ubiquitin chains type, especially in k-48 ubiquitin chain. OTUB1 performs its deubiquitinating function via two distinct manners: the canonical and non-canonical manner. OTUB1 is involved in many critical immune response processes, containing CD8 T cell activation, PD-L1 related immune evasion, and several infectious diseases. OTUB1 plays an important role in the occurrence, proliferation, invasion, and progression of various cancers, such as bladder cancer, breast cancer, prostate cancer, and colorectal cancer (PMID: 34875341, 35296795). OTUB1 has 2 isoforms with molecular weights of 31 and 35 kDa, respectively.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68489-1-Ig (OTUB1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.