

For Research Use Only

STT3B Monoclonal antibody

Catalog Number: 68496-1-Ig



Basic Information

Catalog Number:

68496-1-Ig

Size:

1000 ug/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG30866

GenBank Accession Number:

BC015880

GeneID (NCBI):

201595

UNIPROT ID:

Q8TCJ2

Full Name:

STT3, subunit of the oligosaccharyltransferase complex, homolog B (*S. cerevisiae*)

Calculated MW:

97 kDa

Observed MW:

94 kDa

Purification Method:

Protein G purification

CloneNo.:

3B7A8

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human

Positive Controls:

WB : A549 cells, HeLa cells, HepG2 cells, Jurkat cells, K-562 cells

Background Information

STT3B is a component of the N-oligosaccharyl transferase (OST) enzyme which catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). STT3B seems to be involved in complex substrate specificity

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68496-1-Ig (STT3B antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 68496-1-Ig (STT3B antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



1x10⁶ A431 cells were intracellularly stained with 0.8 ug STT3B Monoclonal antibody (68496-1-Ig, Clone:3B7A8) and Multi-rAb CoraLite[®] Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002)(red), or 0.8 ug Mouse IgG1 isotype control Mouse McAb (66360-1-Ig, Clone: 1F8D3) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).