

For Research Use Only

SMAD7 Monoclonal antibody

Catalog Number: 68594-1-Ig



Basic Information

Catalog Number: 68594-1-Ig	GenBank Accession Number: BC074819	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 4092	CloneNo.: 3D1C3
Source: Mouse	UNIPROT ID: O15105	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG2a	Full Name: SMAD family member 7	
Immunogen Catalog Number: AG13688	Calculated MW: 426 aa, 46 kDa	
	Observed MW: 45-55 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : pig kidney tissue, mouse brain tissue, rat brain tissue, rabbit brain tissue
Species Specificity: Human, mouse, rat, rabbit, pig	

Background Information

SMAD7, also named as Mothers against decapentaplegic homolog 7, is a 426 amino acid protein, which belongs to the dwarfin/SMAD family. SMAD7 Interaction with NEDD4L or RNF111 induces translocation from the nucleus to the cytoplasm (PubMed:16601693). TGF-beta stimulates its translocation from the nucleus to the cytoplasm. PDPK1 inhibits its translocation from the nucleus to the cytoplasm in response to TGF-beta (PubMed:17327236). SMAD7 as antagonist of signaling by TGF-beta type 1 receptor superfamily members has been shown to inhibit TGF-beta and activin signaling by associating with their receptors thus preventing SMAD2 access. SMAD7 functions as an adapter to recruit SMURF2 to the TGF-beta receptor complex and also acts by recruiting the PPP1R15A-PP1 complex to TGFBR1, which promotes its dephosphorylation. SMAD7 positively regulates PDPK1 kinase activity by stimulating its dissociation from the 14-3-3 protein YWHAQ which acts as a negative regulator.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68594-1-Ig (SMAD7 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.