

For Research Use Only

CEACAM1/CD66a Monoclonal antibody

Catalog Number: 68653-1-Ig



Basic Information

Catalog Number:

68653-1-Ig

Size:

1000 ug/ml

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

BC014473

GeneID (NCBI):

634

UNIPROT ID:

P13688

Full Name:

carcinoembryonic antigen-related
cell adhesion molecule 1 (biliary
glycoprotein)

Calculated MW:

526 aa, 58 kDa

Observed MW:

120-150 kDa

Purification Method:

Protein G purification

CloneNo.:

1D1H1

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:500-1:2000

Applications

Tested Applications:

WB, IF/ICC, FC, ELISA

Species Specificity:

human

Positive Controls:

WB : HepG2 cells, HT-29 cells

IF/ICC : HT-29 cells,

Background Information

CEACAM1, also known as CD66a or BGP (biliary glycoprotein), is a member of the carcinoembryonic antigen (CEA) family which belongs to the immunoglobulin superfamily of cell adhesion molecules (PMID: 16919437). CEACAMs play major roles in cell-cell and cell-ECM adhesion and have been implicated in the control of cell proliferation, angiogenesis, and tissue remodeling (PMID: 23021083). CEACAM1 is a transmembrane glycoprotein that is expressed on epithelial, endothelial, and immune cells (PMID: 30314283). CEACAM1 has been demonstrated to play a role in morphogenesis, apoptosis, angiogenesis, cell proliferation, cell motility, fibrosis, and immunity (PMID: 31604530).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68653-1-Ig (CEACAM1/CD66a antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of unfixed HT-29 cells using CEACAM1/CD66a antibody (68653-1-Ig, Clone: 1D1H1) at dilution of 1:1000 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).



1x10⁶ HT-29 cells were surface stained with 0.2 μ g CEACAM1/CD66a Monoclonal antibody (68653-1-Ig, Clone:1D1H1) and CoraLite488-conjugated Goat Anti-Mouse IgG(H+L) (Cat.NO. SA00013-1). Mouse IgG1 isotype control (66360-1-Ig, Clone: 1F8D3) was parallel stained as control. Cells were not fixed.