

For Research Use Only

MUC20 Monoclonal antibody, PBS Only

Catalog Number: 68714-1-PBS



Basic Information

Catalog Number: 68714-1-PBS	GenBank Accession Number: BC029267	Purification Method: Protein A purification
Size: 1mg/ml	GeneID (NCBI): 200958	CloneNo.: 1G9B2
Source: Mouse	UNIPROT ID: Q8N307	
Isotype: IgG3	Full Name: mucin 20, cell surface associated	
Immunogen Catalog Number: AG31840	Calculated MW: 55 kDa, 71 kDa	
	Observed MW: 70 kDa	

Applications

Tested Applications:
WB, IF/ICC, Indirect ELISA

Species Specificity:
human, canine

Background Information

MUC20 contains serine, threonine, and proline-rich repeats in its extracellular domain and is likely to be extensively O-glycosylated. MUC20 blocks GRB2 recruitment to MET thus suppressing the GRB2-RAS pathway and inhibiting HGF-induced proliferation of MMP1 and MMP9 expression (PMID: 15314156). MUC20 is highly expressed in the kidney and localized in the proximal tubules but not in the glomerulus or distal tubules (PMID: 14565953). MUC20 was detected in most of the male urogenital tract epithelia, except for epididymis (PMID: 16997931). The expression of MUC20 is up-regulated in the kidneys of patients with immunoglobulin A nephropathy, in an animal model of lupus nephritis, and in mice with acute renal injury caused by cisplatin administration or unilateral ureteral obstruction (PMID: 14565953).

Storage

Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 68714-1-Ig (MUC20 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 68714-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using MUC20 antibody (68714-1-Ig, Clone: 1G9B2) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 68714-1-PBS in a different storage buffer formulation.