For Research Use Only

ENPP2 Monoclonal antibody

Catalog Number: 68724-1-Ig



Basic Information

Catalog Number:

GenBank Accession Number: BC034961

Purification Method:

68724-1-lg

GeneID (NCBI):

Protein A purification

Size: Source:

CloneNo.:

1000 µg/ml

5168 **UNIPROT ID:** 2A10D9

Mouse Isotype: Q13822

Recommended Dilutions: WB 1:1000-1:4000

IgG2a

AG5499

Full Name: ectonucleotide

Immunogen Catalog Number:

pyrophosphatase/phosphodiesterase

Calculated MW:

99 kDa

Observed MW:

100-110 kDa

Applications

Tested Applications:

Species Specificity:

WB. ELISA

Human, mouse, rat, rabbit, pig

Positive Controls:

WB: pig brain tissue, HuH-7 cells, U-87 MG cells, rabbit brain tissue, rat brain tissue, mouse brain tissue

Background Information

ENPP2, Autotaxin or Extracellular lysophospholipase D, is a 863 amino acid secreted protein, which is detected in blood plasma (at protein level) (PubMed:12176993, PubMed:26371182). ENPP2 is Predominantly expressed in brain, placenta, ovary, and small intestine. ENPP2 is expressed in a number of carcinomas such as hepatocellular and prostate carcinoma, neuroblastoma and non-small-cell lung cancer. ENPP2 is expressed in body fluids such as plasma, cerebral spinal fluid (CSF), saliva, follicular and amniotic fluids. The calculated molecular weight of ENPP2 is about 100 kDa (PMID: 22301782), but modified ENPP2 is 120 kDa (PMID: 19808661). ENPP2 Hydrolyzes lysophospholipids to produce the signaling molecule lysophosphatidic acid (LPA) in extracellular fluids (PubMed:15769751, PubMed:26371182, PubMed:27754931).


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



HuH-7 cells were subjected to SDS PAGE followed by western blot with 68724-1-1g (ENPP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



U-87 MG cells were subjected to SDS PAGE followed by western blot with 68724-1-1g (ENPP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68724-1-1g (ENPP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.