For Research Use Only

ENPP2 Monoclonal antibody

Catalog Number:68724-1-lg 1 Publications



Basic Information

Catalog Number: 68724-1-Ig

Size: 1000 ug/ml Source: Mouse

IgG2a Immunogen Catalog Number:

Immunogen Catalog Number:

AG5499

Isotype:

GenBank Accession Number:

BC034961 GeneID (NCBI): 5168 UNIPROT ID:

Q13822 Full Name: ectonucleotide

pyrophosphatase/phosphodiesterase

2

Calculated MW: 99 kDa Observed MW: 100-110 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat, pig, rabbit

Cited Species:

rat

Positive Controls:

WB: pig brain tissue, HuH-7 cells, U-87 MG cells, rabbit brain tissue, rat brain tissue, mouse brain tissue

Purification Method:

CloneNo.:

2A10D9

Protein A purification

Recommended Dilutions:

WB 1:1000-1:4000

Background Information

ENPP2, Autotaxin or Extracellular lysophospholipase D, is a 863 amino acid secreted protein, which is detected in blood plasma (at protein level) (PubMed:12176993, PubMed:26371182). ENPP2 is Predominantly expressed in brain, placenta, ovary, and small intestine. ENPP2 is expressed in a number of carcinomas such as hepatocellular and prostate carcinoma, neuroblastoma and non-small-cell lung cancer. ENPP2 is expressed in body fluids such as plasma, cerebral spinal fluid (CSF), saliva, follicular and amniotic fluids. The calculated molecular weight of ENPP2 is about 100 kDa (PMID: 22301782), but modified ENPP2 is 120 kDa (PMID: 19808661). ENPP2 Hydrolyzes lysophospholipids to produce the signaling molecule lysophosphatidic acid (LPA) in extracellular fluids (PubMed:15769751, PubMed:26371182, PubMed:27754931).

Notable Publications

Author	Pubmed ID	Journal	Application
Anfal F Bin Dayel	39403839	J Pharm Pharmacol	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



HuH-7 cells were subjected to SDS PAGE followed by western blot with 68724-1-1g (ENPP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



U-87 MG cells were subjected to SDS PAGE followed by western blot with 68724-1-1g (ENPP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68724-1-1g (ENPP2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.