For Research Use Only

NeutraKine® IFN-gamma Mouse McAb Catalog Number:69007-1-lg 1 Publications


proteintech® Antibodies | ELISA kits | Proteins www.ptglab.com

Basic Information	Catalog Number: 69007-1-1g Concentration:	GenBank Accession Number: GeneID (NCBI): 3458		Purification Method: Protein G purification CloneNo.:		
	Source: Mouse	ENSEMBL Gene	ENSEMBL Gene ID: ENSG00000111537		1E10G7	
	Isotype: IgG1 Immunogen Catalog Number: HZ-1301	Full Name: IFN gamma				
Applications	Tested Applications: Neutralization, ELISA Cited Applications: Neutralization Species Specificity: human	Positive Controls: Neutralization : TNF alpha treated HT-29 cells, HT-29 cell ELISA : Recombinant protein,				
Background Information	Interferon gamma (IFNG) is a soluble cytokine that is the only member of the type II class of interferons. It is secreted by Th1 cells, cytotoxic T cells and NK cells. The cytokine is associated with antiviral, immunoregulatory and anti-tumor properties and is a potent activator of macrophages. It plays crucial roles in pathogen clearance. Aberrant IFNG expression is associated with a number of autoinflammatory and autoimmune diseases. It has bee identified in many studies as a biomarker for pleural tuberculosis (TB). Mutations in this gene are associated with aplastic anemia.					
	This antibody can be used to neutralize the bioactivity of Interferon gamma.					
Notable Publications	Author Guanzhan Liang		Journal Cell Mol Gastroente	rol Hepatol	Application Neutralization	
Storage	Storage: Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at (4°C) for short term or at (-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products. Storage Buffer: Sterile PBS, pH7.4 Aliquoting is unnecessary for -20°C storage					


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Recombinant human IFN-gamma (Cat.NO. HZ-1301) inhibits HT-29 cell line (human colorectal adenocarcinoma cell line) proliferation in the prescence of 0.5 ng/mL TNF alpha in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IFN-gamma (2ng/mL HZ-1301 axis) is neutralized by mouse antihuman IFN-gamma monoclonal antibody 69007-1-lg at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 100-



Recombinant human IFN-gamma (Cat.NO. HZ-1301) inhibits HT-29 cell line (human colorectal adenocarcinoma cell line) proliferation in the prescence of 0.5 ng/mL TNF alpha in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IFN-gamma (5 ng/mL HZ-1301 axis) is neutralized by mouse antihuman IFN-gamma monoclonal antibody 69007-1-lg at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 100-



Indirect ELISA was carried out by coating recombinant Human IFN gamma (Cat.NO. HZ-1301) at 70 ng/well followed by blocking and adding serial diluted IFN gamma antibody 69007-1-Ig and 69507-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.