For Research Use Only

NeutraKine® BMP-7 Monoclonal antibody



Catalog Number: 69011-1-Ig

Basic Information

Catalog Number:

69011-1-lg

Size: Source:

Mouse

Isotype:

igui

Immunogen Catalog Number:

HZ-1229

GenBank Accession Number:

GeneID (NCBI):

655 Full Name:

bone morphogenetic protein 7

Purification Method: Protein G purification

CloneNo.: 3E9G7

Applications

Tested Applications:

FC (Intra), Neutralization, ELISA

Species Specificity:

human, mouse

Background Information

The bone morphogenetic proteins (BMPs) are a family of secreted signaling molecules that can induce ectopic bone growth. Many BMPs are part of the transforming growth factor-beta (TGFB) superfamily. BMPs were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extrackeletal site.

BMP7, also known as osteogenic protein-1 or OP-1, plays a key role in the transformation of mesenchymal cells into bone and cartilage. BMP7 may be involved in bone homeostasis (PMID: 15621726). It is expressed in the brain, kidneys and bladder. BMP7 is also present in cancers, including breast, prostate, and colon cancers, in which it is implicated in regulating cancer cell proliferation (PMID: 16419056, PMID: 15531927). Overexpression of BMP7 mRNA in colorectal cancer patients was significantly associated with poor prognosis and low overall survival (PMID: 18259822). Recent studies suggest that high-expression level of BMP7 serves as a biomarker for poor prognosis for HCC (PMID: 23179403).

This antibody can be used to neutralize the bioactivity of BMP-7.

Storage

Storage


Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at(4°C) for short term or at(-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.


Storage Buffer:

Sterile PRS

Aliquoting is unnecessary for -20°C storage

Selected Validation Data





Recombinant human BMP-7 (Cat.NO. HZ-1229) induces alkaline phosphatase production in the ATDC-5 cell line (using pNPP as chromogenic substrate for detection) in a dose dependent manner (blue curve, bottom X-left Y). The activity of human BMP-7 (200 ng/mL HZ-1229) is neutralized by mouse anti-human BMP-7 antibody 69011-1-lg at serial dose (red curve, refer to top X-right Y). The ND50 is typically 1-2 µ g/mL

1X10^6 HEK-293 cells were intracellularly stained with 0.4 ug Anti-Human NeutraKine® BMP-7 (69011-1-lg, Clone:3E9G7) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).