For Research Use Only

NeutraKine® SCF Monoclonal antibody




Catalog Number:69026-1-lg

Basic Information	Catalog Number: 69026-1-1g Size: 1000 µ g/ml Source: Mouse Isotype: IgG1 Immunogen Catalog Number: HZ-1024	GenBank Accession Number: GeneID (NCBI): 4254 Full Name: KIT ligand	Purification Method: Protein G purification CloneNo.: 1B3C4
Applications	Tested Applications: Neutralization, ELISA Species Specificity: Human		
Background Information	Stem cell factor (SCF) also known as mast cell growth factor, steel factor, and kit ligand, is a hematopoietic growth factor. SCF is a critical protein with key roles in the cell such as hematopoiesis, gametogenesis and melanogenesis. SCF activates multiple signal pathways by binding with c-kit receptor. SCF and its receptor, c-kit ligand (KL), are up-regulated in particular human malignancies including gastrointestinal stromal tumor (GISTs), breast cancer, hematopoietic cell, myeloid leukaemia, and glioma. The SCF/c-KIT system also plays a relevant role in cell fate decision, and its activity is widely known to control cell proliferation, differentiation, survival and apoptosis. This antibody is used to neutralize the bioactivity of SCF.		
Storage	reconstitution we recommend that t	he solution can be stored at(4°C) for sl hould be avoided with reconstituted p	red between (-20°C) and (-80°C). Upon hort term or at(-20°C) to (-80°C) for long rroducts.

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human SCF (Cat.NO. HZ-1024) stimulates proliferation of TF-1 cell line (human erythroleukemic cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human SCF (20 ng/mL HZ-1024) is neutralized by mouse anti-human SCF monoclonal antibody 69026-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 0.5-2 μ g/mL