

For Research Use Only

NeutraKine®HGF Monoclonal antibody, PBS Only

Catalog Number: 69027-1-PBS



Basic Information

Catalog Number:

69027-1-PBS

Concentration:

1 mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

HZ-1084

GenBank Accession Number:

GeneID (NCBI):

3082

Full Name:

hepatocyte growth factor
(hepapoietin A; scatter factor)

Purification Method:

Protein G purification

CloneNo.:

2D2B4

Applications

Tested Applications:

Neutralization, ELISA

Species Specificity:

human

Background Information

Hepatocyte growth factor (HGF) is the most potent mitogen of mature hepatocytes in primary culture. HGF is derived from a biologically inactive single chain precursor of 728 amino acids (pro-HGF) localized mostly on the cell surface and in the extracellular matrix. The mature form produced following proteolytic cleavage is composed of a 69-kDa α -subunit (containing four kringle domains) and the 34 kDa β -subunit, similar to the catalytic domain of serine proteases, but with amino acid substitutions in the active site. HGF is a pleiotropic cytokine which exerts a variety of effects on several cells, being involved in the regulation of many biological processes, such as inflammation, tissue repair, morphogenesis, angiogenesis, tumour propagation, immunomodulation of viral infections and cardio-metabolic activities.

This antibody is used to neutralize the bioactivity of HGF.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only, pH7.4

For technical support and original validation data for this product please contact:



T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human HGF (Cat.NO. HZ-1084) stimulates Saos-2 cells (human osteosarcoma cell) produce IL-11 in a dose dependent manner (blue curve, refer to bottom X-Left Y axis). The activity of human HGF (50 ng/mL HZ-1084) is neutralized by mouse anti-human HGF monoclonal antibody 69027-1-Ig (red curve, refer to top X-right Y axis). The ND50 is typically 2-8 μ g/mL. This data was developed using the same antibody clone with 69027-1-PBS in a different storage buffer formulation.

Indirect ELISA was carried out by coating recombinant Human HGF (Cat.NO. HZ-1084) at 70 ng/well followed by blocking and adding serial diluted Thrombopoietin antibody 69027-1-Ig and 69527-1-Ig respectively. Signal was developed with TMB and stopped by H₂SO₄. Signal strength was measured by absorbance at 450 nm. This data was developed using the same antibody clone with 69027-1-PBS in a different storage buffer formulation.