

For Research Use Only

NeutraKine®IL-27 Monoclonal antibody



Catalog Number: 69034-1-Ig

Basic Information

Catalog Number:

69034-1-Ig

Size:

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

HZ-1275

GenBank Accession Number:

GeneID (NCBI):

246778

UNIPROT ID:

Q8NEV9

Full Name:

interleukin 27

Purification Method:

Protein A purification

CloneNo.:

2H2A2

Applications

Tested Applications:

ELISA, Neutralization

Species Specificity:

Human

Background Information

IL-27 is a heterodimeric cytokine composed of two subunits: IL-27a and IL-27b. IL-27 acts on various cell types, including T cells, B cells, macrophages, dendritic cells, natural killer (NK) cells and non-hematopoietic cells. IL-27 plays a critical role in the early regulation of T helper type 1 initiation, and enhances proliferation of naive CD4+T cells and naive B cells. It, however, also exerts anti-inflammatory functions by inhibiting the development of Th17 cells and inducing IL-10 producing type 1 regulatory T cells. IL-27 is a potentially promising cytokine for therapeutic approaches on various human diseases.

This antibody is used to neutralize the bioactivity of IL-27.

Storage

Storage:

Lyophilized antibodies are stable for 1 year from the date of receipt if stored between (-20°C) and (-80°C). Upon reconstitution we recommend that the solution can be stored at(4°C) for short term or at(-20°C) to (-80°C) for long term. Repeated freeze thaw cycles should be avoided with reconstituted products.

Storage Buffer:

Sterile PBS.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Recombinant human IL-27 (Cat.NO. HZ-1275) stimulates proliferation of OCI-AML5 cells in a dose dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IL-27 (250ng/mL HZ-1275) is neutralized by mouse anti-human IL-27 monoclonal antibody 69034-1-Ig at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 0.4-1 µg/mL.



Indirect ELISA was carried out by coating recombinant Human IL-27 (Cat.NO. HZ-1275) at 70 ng/well followed by blocking and adding serial diluted IL-27 antibody 69034-1-Ig and 69534-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.