

For Research Use Only

FITC Recombinant antibody

Catalog Number: 80003-1-RR **1 Publications**



Basic Information

Catalog Number:

80003-1-RR

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

GeneID (NCBI):

Full Name:

Purification Method:

Protein A purification

CloneNo.:

1B2

Recommended Dilutions:

WB 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

IP

Species Specificity:

FITC conjugated

Cited Species:

human

Positive Controls:

WB : FITC conjugated antibody,

Background Information

Fluorescein isothiocyanate (FITC) is a derivative of fluorescein used in wide-ranging applications. It is the original fluorescein molecule functionalized with an isothiocyanate reactive group (-N=C=S), replacing a hydrogen atom on the bottom ring of the structure. It is typically available as a mixture of isomers, fluorescein 5-isothiocyanate (5-FITC) and fluorescein 6-isothiocyanate (6-FITC). FITC is reactive towards nucleophiles including amine and sulfhydryl groups on proteins. CAS number of FITC: 27072-45-3 (mixture); 3326-32-7, 5-isomer; 18861-78-4 (6-isomer). This rabbit recombinant antibody detects the FITC only. This antibody can be coated in the plate to capture the FITC conjugated protein or antibody.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|--------------|-----------|---------------------|-------------|
| Jialiang Luo | 36029544 | Biomed Pharmacother | IP |

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



FITC-conjugated secondary antibodies were subjected to SDS PAGE followed by western blot with 80003-1-RR (FITC antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.