

For Research Use Only

SARS-CoV-2 Nucleocapsid Phosphoprotein Recombinant antibody

Catalog Number: 80026-1-RR



Basic Information

Catalog Number:

80026-1-RR

Size:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG30676

GenBank Accession Number:

NC_045512

GeneID (NCBI):

43740575

Full Name:

COVID-19 N Protein

Purification Method:

Protein A purification

CloneNo.:

4B9

Recommended Dilutions:

WB 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

virus, recombinant protein

Positive Controls:

WB : Eukaryotic nucleocapsid phosphoprotein,
Recombinant protein

Background Information

The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. It can be used as a marker in diagnostic assays due to its high immunogenicity (PMID: 32416961, PMID: 32235387). A sandwich ELISA for COVID-19 N Protein can be assembled by using 80027-1-RR as capture antibody and conjugated 80026-1-RR for detection.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




E.coli expressed SARS-CoV-2 Nucleocapsid Phosphoprotein (Cat.NO. Ag30676) was subjected to SDS-PAGE followed by western blot with 80027-1-RR and 80026-1-RR at various work concentration.



Eukaryotic expressed SARS-CoV-2 Nucleocapsid Phosphoprotein was subjected to SDS-PAGE followed by western blot with 80027-1-RR and 80026-1-RR at various work concentration.



Indirect ELISA was carried out by coating eukaryotic expressed N protein at 70 ng/well followed by blocking and adding serial diluted primary antibody 80026-1-RR and 80027-1-RR respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.



Sandwich ELISA was carried out by coating 80027-1-RR at 80 ng/well followed by blocking and adding different concentration of eukaryotic expressed N protein (0.5-1000 ng/mL). HRP-conjugated 80026-1-RR was used at 1 µg/mL for detection. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.