

For Research Use Only

Phospho-Beta Catenin (Ser552) Recombinant antibody

Catalog Number: 80035-2-RR



Basic Information

Catalog Number:

80035-2-RR

Concentration:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC058926

GeneID (NCBI):

1499

ENSEMBL Gene ID:

ENSG00000168036

UNIPROT ID:

P35222

Full Name:

catenin (cadherin-associated protein),
beta 1, 88kDa

Calculated MW:

781 aa, 86 kDa

Observed MW:

86-100 kDa

Purification Method:

Protein A purification

CloneNo.:

250272E1

Recommended Dilutions:

WB: 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human

Positive Controls:

WB : Forskolin treated HeLa cells, Forskolin treated
A431 cells

Background Information

β-Catenin, also known as CTNNB1, is an evolutionarily conserved, multifunctional intracellular protein. β-Catenin is a 92-kDa protein, originally identified in cell adherens junctions (AJs) where it functions to bridge the cytoplasmic domain of cadherins to α-catenin and the actin cytoskeleton. Besides its essential role in the AJs, β-catenin is also a key downstream component of the canonical Wnt pathway that plays diverse and critical roles in embryonic development and adult tissue homeostasis. The Wnt/β-catenin pathway is also involved in the activation of other intracellular messengers such as calcium fluxes, JNK, and SRC kinases. Deregulation of β-catenin activity is associated with multiple diseases including cancers (PMID: 22617422, PMID: 18334222). Both AMPK and AKT phosphorylates β-catenin at Ser 552. The phosphorylation of β-catenin at Ser552 can stabilize β-catenin, enhance its nuclear accumulation and increase its transcriptional activity (PMID: 2865606, PMID: 20361929).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated A431 cells and Forskolin treated A431 cells were subjected to SDS PAGE followed by western blot with 80035-2-RR (Phospho-Beta Catenin (Ser552) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control.



Non-treated HeLa cells and Forskolin treated HeLa cells were subjected to SDS PAGE followed by western blot with 80035-2-RR (Phospho-Beta Catenin (Ser552) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control.