

For Research Use Only

Phospho-STK11/LKB1 (Thr189) Recombinant antibody



Catalog Number: 80127-1-RR **1 Publications**

Basic Information

Catalog Number: 80127-1-RR	GenBank Accession Number: BC007981	Purification Method: Protein A purification
Size: 250 µg/ml	GeneID (NCBI): 6794	CloneNo.: 5J9
Source: Rabbit	UNIPROT ID: Q15831	Recommended Dilutions: WB 1:1000-1:10000 IF 1:50-1:500
Isotype: IgG	Full Name: serine/threonine kinase 11	
	Calculated MW: 49 kDa	
	Observed MW: 50-55 kDa	

Applications

Tested Applications: FC, IF/ICC, WB, ELISA	Positive Controls: WB : PC-3 cells, HEK-293 cells, NIH/3T3 cells, Calyculin A treated HEK-293 cells, Calyculin A treated PC-3 cells, Calyculin A treated NIH/3T3 cells
Cited Applications: WB	IF : Calyculin A treated PC-3 cells,
Species Specificity: Human, Mouse	
Cited Species: mouse	

Background Information

STK11 (serine/threonine-protein kinase 11) is also named as LKB1, PJS, and belongs to the protein kinase superfamily. It controls the activity of AMP-activated protein kinase (AMPK) family members, thereby playing a role in various processes such as cell metabolism, cell polarity, apoptosis and DNA damage response. The tumour suppressor protein LKB1 is a serine/threonine kinase that has been causally linked to Peutz-Jeghers syndrome (PJS). Defects in STK11 are a cause of Peutz-Jeghers syndrome (PJS) and defects in STK11 have been associated with testicular germ cell tumor (TGCT) and some sporadic cancers, especially lung cancers.

Notable Publications

Author	Pubmed ID	Journal	Application
Bingbing Liu	38452099	PLoS One	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Non-treated PC-3 and Calyculin A treated PC-3 cells were subjected to SDS PAGE followed by western blot with 80127-1-RR (Phospho-STK11/LKB1 (Thr189) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Immunofluorescent analysis of (4% PFA) fixed non-treated PC-3 cells and Calyculin A treated PC-3 cells using Phospho-STK11/LKB1 (Thr189) antibody (80127-1-RR, Clone: 5J9) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10⁶ PC-3 cells untreated (dashed lines) or Calyculin A treated (red) were intracellularly stained with 0.25 ug Anti-Human Phospho-STK11 (Thr189) (80127-1-RR, Clone:5J9) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.25 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.