For Research Use Only

STAT3 Recombinant antibody

Catalog Number:80149-1-RR

Featured Product

2 Publications



Purification Method:

CloneNo.:

Protein A purification

Recommended Dilutions:

WB 1:1000-1:8000

Basic Information

Catalog Number: 80149-1-RR

Concentration: 400 ug/ml Source:

Rabbit Isotype:

IgG

GenBank Accession Number:

BC000627 GeneID (NCBI): 6774 UNIPROT ID:

P40763 Full Name:

signal transducer and activator of transcription 3 (acute-phase response factor)

Calculated MW: 770 aa, 88 kDa Observed MW:

88 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity: human, mouse Cited Species: human, mouse Positive Controls:

WB: PC-12 cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, NIH/3T3 cells, Neuro-2a cells

Background Information

Signal transducer and activator of transcription 3 (acute-phase response factor) (STAT3, synonyms: APRF, FLI20882, MGC 16063) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT3 is activated through phosphorylation in response to various cytokines and growth factors including IFNs, EGF, IL5, IL6, HGF, LIF and BMP2. STAT3 mediates the expression of a variety of genes in response to cell stimuli, and thus plays a key role in many cellular processes such as cell growth and apoptosis. The small GTPase Rac1 has been shown to bind and regulate the activity of STAT3.

Notable Publications

Author	Pubmed ID	Journal	Application
Ting Li	39575303	MedComm (2020)	WB
Xiaoning Wei	37952482	Int Immunopharmacol	WB

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 80149-1-RR (STAT3 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 80149-1-RR (STAT3 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



WB result of STAT3 antibody (80149-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STAT3 transfected HeLa cells.