For Research Use Only

## PEX14 (human specific) Recombinant antibody

Catalog Number:80196-1-RR



**Basic Information** 

Catalog Number: GenBank Accession Number: 80196-1-RR BC006327

 Concentration:
 GeneID (NCBI):

 700 ug/ml
 5195

 Source:
 UNIPROT ID:

Rabbit 075381 Isotype: Full Name:

IgG peroxisomal biogenesis factor 14

Immunogen Catalog Number: Calculated MW:

AG0932 41 kDa

Observed MW:

57 kDa

Purification Method: Protein A purification

CloneNo.:

6K5 Recommended Dilutions: WB 1:2000-1:14000

IHC 1:200-1:800 IF/ICC 1:50-1:500

Applications

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

**Positive Controls:** 

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat

cells

IHC: human liver cancer tissue, human kidney tissue

IF/ICC: HeLa cells,

**Background Information** 

PEX14 (peroxisomal biogenesis factor 14) is a peroxisomal membrane protein that is essential for protein docking onto the peroxisomes. It is a central component of the peroxisimal matrix protein import machinery and interacts with PEX5 and PEX19. PEX14 is ubiquitously expressed and defects in PEX14 are the cause of peroxisome biogenesis disorder complementation group K (PBD-CGK). This antibody can be used to detect endogenous PEX14 with an apparent molecular weight of 57 kDa (PMID: 16449325; 9653144) and recognize peroxisomal structures in human, monkey and mouse cells.

This antibody is specific to human PEX14. It can't detect mouse or rat PEX14.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 80196-1-RR (PEX14 (human specific) antibody) at dilution of 1:7000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 80196-1-RR (PEX14 (human specific) antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 80196-1-RR (PEX14 (human specific) antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using PEX14 (human specific) antibody (80196-1-RR, Clone: 6K5 ) at dilution of 1:200 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human PEX14 (human specific) (80196-1-RR, Clone:6K5) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 80196-1-RR against PEX14 (human specific) were performed. The affinity constant is 0.613 nM.