For Research Use Only

## Phospho-S6 Ribosomal protein (Ser236) Recombinant antibody

Catalog Number:80206-1-RR 1 Publications



**Basic Information** 

Catalog Number: 80206-1-RR Concentration:

250 µg/ml

Rabbit Isotype:

IgG

GenBank Accession Number:

BC000524 GeneID (NCBI): 6194

UNIPROT ID: P62753 Full Name:

ribosomal protein S6 Calculated MW: 29 kDa

Observed MW: 32 kDa

Purification Method:

Protein A purification

CloneNo.: 7K17

Recommended Dilutions:

WB 1:2000-1:10000

**Applications** 

Tested Applications: WB, FC (Intra), ELISA Cited Applications:

WB

Species Specificity: human, mouse Cited Species: mouse **Positive Controls:** 

WB: HEK-293 cells, MCF-7 cells, Calyculin A treated

HEK-293 cells, IGF-1 treated MCF-7 cells

## **Background Information**

Ribosomal protein S6 (RPS6) is one of the components of the 40S ribosomal subunit. RPS6 has been functionally regarded as the stimulator and/or inhibitor of certain types of mRNA translation, as well as the regulator of cellular metabolisms, cells size, survival and proliferation. RPS6 is phosphorylated at multiple sites, comprised between Ser235 and Ser247, by the p70 rpS6 kinase (S6K) 1, which is a major downstream effector of the mammalian target of rapamycin complex 1 (mTORC1). Phosphorylation of RPS6 at the dual site Ser235/236 occurs also independently of mTORC1, via the p90 ribosomal S6 kinases (RSK), which are activated by the extracellular signal-regulated kinases (ERK). Recent studies performed in pancreatic  $\beta$ -cells identified PKA as an additional RPS6 kinase, specifically involved in the phosphorylation of Ser235/236. (PMID: 26490682, PMID: 21814187, PMID: 31112404). 80206-1-RR specifically recognizes the phosphorylation site of Ser236 or dual site Ser235/236.

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Jie Li	39702426	NPJ Biofilms Microbiomes	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**



Non-treated and Calyculin A treated HEK-293 cells were subjected to SDS PAGE followed by western blot with 80206-1-RR (Phospho-56 Ribosomal protein (Ser236) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



1X10^6 HEK-293 cells untreated (dashed lines) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug Anti-Human Phospho-S6 Ribosomal protein (Ser236) (80206-1-RR, Clone: PK.17) and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000, or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.