For Research Use Only

P62,SQSTM1 Recombinant antibody, **PBS Only**

Catalog Number:80294-1-PBS

Featured Product



Purification Method:

Protein A purification

CloneNo.:

1D17

Basic Information

Catalog Number: 80294-1-PBS Concentration: 1 mg/ml

Source: Rabbit Isotype: Immunogen Catalog Number:

AG13131 48 kDa

> Observed MW: 62 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), IP, Indirect ELISA

Species Specificity:

human

Background Information

Sequestosome 1 (SQSTM1/p62) is a multifunctional adaptor protein implicated in selective autophagy, cell signaling pathways, and tumorigenesis. p62 has been implicated in shuttling ubiquitinated and aggregated proteins for autophagic degradation. p62 is degraded during the autophagic process, which makes its intracellular level a $marker for autophagy\ progression.\ p62\ is\ at\ the\ cross-roads\ of\ several\ signaling\ pathways\ including\ Ras/\ Raf/\ MAPK$ and NF \upkepsilon B and plays important role in cancer. p62 is a component of inclusion bodies/ protein aggregates found in human diseases, including Huntington's disease, Alzheimer's disease, Parkinson's disease, and nephropathic cystinosis (PMID: 22074114, 22860231, 22714671).

GenBank Accession Number:

BC017222

8878

Q13501 Full Name:

GeneID (NCBI):

UNIPROT ID:

sequestosome 1 Calculated MW:

Storage


Storage:


Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only

Selected Validation Data










Lysates of HepG2 cells treated with Chloroquine or not were subjected to SDS PAGE followed by western blot with 80294-1-RR (P62,SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

WB result of P62,SQSTM1 antibody (80294-1-RR; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-P62,SQSTM1 transfected HEK-293 cells. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

Various lysates were subjected to SDS PAGE followed by western blot with 80294-1-RR (P62,SQSTM1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.








Immunofluorescent analysis of (-20°C Ethanol) fixed Chloroquine treated HeLa cells using P62,SQSTM1 antibody (80294-1-RR, Clone: 1D17) at dilution of 1:500 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

1X10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human P62,5Q5TM1 (80294-1-RR, Clone:1D17) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.

IP result of anti-P62,SQSTM1 (IP:80294-1-RR, 4ug; Detection:80294-1-RR 1:500) with HEK-293 cells lysate 1640 ug. This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human pancreas cancer tissue slide using 80294-1-RR (P62,SQSTM1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80294-1-PBS in a different storage buffer formulation