For Research Use Only

Phospho-Hamartin/TSC1 (Ser511) Recombinant antibody

Catalog Number:80340-1-RR 1 Publications





Basic Information	Catalog Number: 80340-1-RR	GenBank Accession Number: NM_000368		Purification Method: Protein A purification		
	Size: 500 µg/ml	GenelD (NCBI): 7248		CloneNo.: 3N6		
	Source:UNIPROT ID:RabbitQ92574			Recommended Dilutions: WB 1:5000-1:50000		
	lsotype: IgG	Full Name: tuberous sclerosis :	1			
		Calculated MW: 130 kDa				
		Observed MW: 150 kDa				
Applications	Tested Applications:		Positive C	Positive Controls:		
	WB, FC (Intra), ELISA Cited Applications:		HeLa cells	WB: NIH/3T3 cells, HeLa cells, Calyculin A treated HeLa cells, Calyculin A treated HEK-293 cells,		
	WB, IF Calyculin A treated NIH/3T3 cells Species Specificity: human, mouse					
	Cited Species: mouse					
Background Information	TSC1, also named as Hamartin, is a tumor suppressor gene syndrome whose manifestations can include seizures, mental retardation, autism, and tumors in the brain, retina, kidney, heart, and skin. The tuberous sclerosis 1 (TSC1)/TSC2 tumor suppressor complex serves as a repressor of the mTOR pathway, and disruption of TSC1/TSC2 complex function may contribute to tumorigenesis. TSC1 is phosphorylated by IKKb at Ser487 and Ser511 in vitro and in vivo, and phosphorylation at these sites enhances dissociation of the TSC1/TSC2 complex, which in turn induces mTOR activation. (PMID: 17693255). TSC1 is also threonine-phosphorylated at three sites (Thr417, Ser584, and Thr1047) in a reaction catalyzed by cyclin-dependent kinase 1 (CDK1) (PMID: 14551205).					
Notable Publications	Author	Pubmed ID Jou	ırnal	Application		
	Xiaoxue Jiang	39261456 Na	t Commun	WB,IF		
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azid Aliquoting is unnecessary fo	le and 50% glycerol pH 7.3.				

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Non-treated and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80340-1-RR (Phospho-Hamartin/TSC 1 (Ser511) Antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. Non-treated and Calyculin A treated HeLa and HEK-293 cells were subjected to SDS PAGE followed by western blot with 80340-1-RR (Phospho-Hamartin/TSC1 (Ser511) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1X10^6 HeLa cells untreated (dashed lines) and Calyculin A treated (red) were intracellularly stained with 0.5 ug Anti-Human Phospho-Hamartin/TSC1 (Ser511) (80340-1-RR, Clone:3N6) and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) at dilution 1:1000, or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 50% MeOH.