For Research Use Only

Perilipin-2 Recombinant antibody

Catalog Number:80362-2-RR



Basic Information

Catalog Number: GenBank Accession Number: 80362-2-RR BC005127

Concentration: GenelD (NCBI): 1000 ug/ml 123

Source: UNIPROT ID:
Rabbit Q99541
Isotype: Full Name:

gG adipose differentiation-related

Immunogen Catalog Number: protein

G7539 Calculated MW:

48 kDa Observed MW: 48 kDa Purification Method:

Protein A purification

CloneNo.: 230274F4

Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500 IF/ICC 1:125-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: THP-1 cells, HepG2 cells, K-562 cells

IHC: mouse liver tissue,

IF/ICC: oleic acid treated HeLa cells,

Background Information

ADRP (adipocyte differentiation related protein) also known as ADFP, adipophilin, or perilipin-2, is a member of PAT family which is responsible for the transportation of lipids and the formation of lipid droplets. ADRP is localized on the surface of lipid droplets in a variety of tissues and cell lines. ADRP is not detected in undifferentiated cells but increases rapidly to high levels when adipocyte precursors differentiate into adipocytes. Anti-ADRP antibody is a reliable and sensitive marker for lipid droplet. Enhanced expression of ADRP is linked to diseases with abnormal lipid storage, including hepatic steatosis, atherosclerosis and diabetes. Immunohistochemistry of ADRP may facilitate histomorphological diagnosis of these diseases.

Storage


Storage:

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80362-2-RR (ADRP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




1x10^6 HepG2 cells were intracellularly stained with 0.25 ug ADRP Recombinant antibody (80362-2-RR, Clone:230274F4) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 80362-2-RR (ADRP antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse liver tissue slide using 80362-2-RR (ADRP antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Various lysates were subjected to SDS PAGE followed by western blot with 80362-2-RR (ADRP antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (-20°C Ethanol) fixed oleic acid treated HeLa cells using ADRP antibody (80362-2-RR, Clone: 230274F4) at dilution of 1:800 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).



Immunofluorescent analysis of (-20°C Ethanol) fixed oleic acid treated Hela cells using ADRP antibody (80362-2-RR, Clone: 230274F4) at dilution of 1:800 and Multi-rAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).



Immunofluorescent analysis of (-20°C Ethanol) fixed oleic acid treated HeLa cells using ADRP antibody (80362-2-RR, Clone: 230274F4) at dilution of 1:250 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLI) kinetic assays of 80362-2-RR against Human ADRP were performed. The affinity constant is below 1 pM.