

For Research Use Only

Phospho-AKT1 (Ser473) Recombinant antibody

Catalog Number: 80462-1-RR

2 Publications



Basic Information

Catalog Number:

80462-1-RR

Concentration:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_005163

GeneID (NCBI):

207

UNIPROT ID:

P31749

Full Name:

v-akt murine thymoma viral
oncogene homolog 1

Observed MW:

56-62 kDa

Purification Method:

Protein A purification

CloneNo.:

2M10

Recommended Dilutions:

WB 1:2000-1:10000

Applications

Tested Applications:

WB, FC (Intra), ELISA

Cited Applications:

WB

Species Specificity:

human, mouse

Cited Species:

human, mouse

Positive Controls:

WB: HeLa cells, HEK-293T cells, HEK-293 cells,
NIH/3T3 cells, IGF-1 treated HEK-293T cells, Calyculin
A treated HEK-293 cells, Calyculin A treated HeLa
cells, Calyculin A treated NIH/3T3 cells

Background Information

AKT is a serine/threonine kinase and it participates in the key role of the PI3K signaling pathway. Phosphatidylinositol-3 kinase (PI3K) is the key regulator of AKT activation. The recruitment of inactive AKT protein to PIP3-rich areas of the plasma membrane results in a conformational change that exposes the activation loop of AKT. AKT's activating kinase, phosphoinositide-dependent protein kinase (PDK1), is also recruited to PIP3 microdomains. PDK1 phosphorylates AKT on threonine 308 (Thr308) of the exposed activation loop, activating AKT and leading to a second phosphorylation of AKT at serine 473 (Ser473) by a kinase presumed to be mTORC2 that further potentiates kinase activity. Active AKT will phosphorylate various downstream protein targets that control cell growth and translational control and act to suppress apoptosis. (PMID: 31594388, PMID: 30808672). 80462-1-RR specifically recognizes AKT1 phosphorylated at Ser473.

Notable Publications

Author	Pubmed ID	Journal	Application
Kun Wang	35415320	ACS Omega	WB
Yihan Huang	38092386	Biol Pharm Bull	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 80462-1-RR (Phospho-AKT (Ser473) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



Non-treated and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 80462-1-RR (Phospho-AKT (Ser473) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with GAPDH antibody as loading control.



1X10⁶ NIH/3T3 cells untreated (dashed line) or treated with Calyculin A (red) were intracellularly stained with 0.5 ug Anti-Human Phospho-AKT1 (Ser473) (80462-1-RR, Clone:2M10) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000, or 0.5 ug Control Antibody (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.