For Research Use Only

TOM20 Recombinant antibody

Catalog Number:80501-1-RR



Basic Information

Catalog Number: GenBank Accession Number: 80501-1-RR BC000882

 Size:
 GeneID (NCBI):

 500 µg/ml
 9804

 Source:
 UNIPROT ID:

 Rabbit
 Q15388

Isotype: Full Name:
IgG translocase of outer mitochondrial

Immunogen Catalog Number: membrane 20 homolog (yeast)

AG2378 Calculated MW: 145 aa, 16 kDa

Observed MW: 16 kDa Purification Method:

Protein A purification CloneNo.:

1D22 Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500

IF 1:50-1:500

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity:

Human, mouse, rat, pig, chicken, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, zebrafish tissue, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, pig brain tissue, rat brain tissue, mouse brain tissue, chicken brain tissue

IHC: human liver tissue,
IF: HepG2 cells,

Background Information

TOM20, also named as KIAA0016, belongs to the Tom20 family. It is a central component of the receptor complex responsible for the recognition and translocation of cytosolically synthesized mitochondrial preproteins. Together with TOM22, TOM20 functions as the transit peptide receptor at the surface of the mitochondrion outer membrane and facilitates the movement of preproteins into the TOM40 translocation pore. TOM20 is characterized as major docking receptors to mediate the recognition by different mechanisms.

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 80501-1-RR (TOM20 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using TOM20 antibody (80501-1-RR, Clone: 1D22) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 80501-1-RR (TOM20 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).