

For Research Use Only

iNOS Recombinant antibody, PBS Only

Catalog Number: 80517-1-PBS



Basic Information

Catalog Number:

80517-1-PBS

Size:

1mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_000625

GeneID (NCBI):

4843

UNIPROT ID:

P35228

Full Name:

nitric oxide synthase 2, inducible

Calculated MW:

131 kDa

Observed MW:

110-130 kDa

Purification Method:

Protein A purification

CloneNo.:

6022

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human, mouse, rat

Background Information

NOS2, also named as iNOS and NOS2A, produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. NO is a reactive free radical which acts as a biologic mediator in several processes, including neurotransmission, antimicrobial and antitumoral activities. NOS2 is a nitric oxide synthase which is expressed in liver and is inducible by a combination of lipopolysaccharide and certain cytokines. iNOS has a very short half-life due to rapid degradation by calpain. iNOS monomer is a direct substrate of calpain I and can be cleaved by calpain I at the canonical CaM-binding site(503-532aa) of iNOS, and then a ~70kD band can be detected by western (PMID:11786228). This antibody is specific to NOS2.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Untreated and LPS treated RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 80517-1-RR (iNOS antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.




LPS treated human PBMCs were subjected to SDS PAGE followed by western blot with 80517-1-RR (iNOS antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 80517-1-RR (iNOS antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 80517-1-PBS in a different storage buffer formulation.