

IRF3 Recombinant antibody

Catalog Number: 80519-1-RR

Basic Information

Catalog Number: 80519-1-RR	GenBank Accession Number: BC009395	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 3661	CloneNo.: 2H6
Source: Rabbit	UNIPROT ID: Q14653	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG	Full Name: interferon regulatory factor 3	IHC 1:50-1:500 IF/ICC 1:50-1:500
	Calculated MW: 47 kDa	
	Observed MW: 50-60 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, ELISA

Species Specificity:
human, mouse, zebrafish

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, THP-1 cells, HT-29 cells, RAW 264.7 cells, zebrafish tissue, HepG2 cells, Jurkat cells

IHC : human breast cancer tissue, human colon tissue

IF/ICC : HeLa cells,

Background Information

The virus-induced expression of interferon (IFN) genes in infected cells implicate in the interplay of several constitutively expressed and virus-activated transcription factors. A family of IFN regulatory factors (IRFs) have been shown to have a role in the transcription of IFN genes as well as IFN-stimulated genes. IRF3 is a novel key transcriptional regulator of type I IFN-dependent immune responses and involves in the innate immune response against DNA and RNA viruses, by binding to the promoters of IFN. It is located in the cytoplasm of uninfected cells in an inactive form, and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, could be phosphorylated by IKKε and TBK1 kinases. This induces a conformational change, leading to its dimerization, nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DAF1), a complex which activates the transcription of the type I IFN and ISG genes.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
 Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80519-1-RR (IRF3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



RAW 264.7 cells were subjected to SDS PAGE followed by western blot with 80519-1-RR (IRF3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 80519-1-RR (IRF3 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using IRF3 antibody (80519-1-RR, Clone: 2H6) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).