For Research Use Only


Phospho-YAP1 (Ser127) Recombinant antibody, PBS Only Catalog Number:80694-2-PBS




Basic Information	Catalog Number: 80694-2-PBS	GenBank Accession Number: BC038235	Purification Method: Protein A purfication
	Concentration: 1 mg/ml	GeneID (NCBI): 10413	CloneNo.: 240779G4
	Source: Rabbit	UNIPROT ID: P46937	
	Isotype: IgG	Full Name: Yes-associated protein 1, 65kDa	
		Calculated MW: 504 aa, 54 kDa	
		Observed MW: 70-78 kDa	
Applications	Tested Applications: WB, FC (Intra), ELISA		
	Species Specificity: human		
Background Information	The transcriptional factors Yes-associated protein 1 (YAP1) and PDZ-binding motif (TAZ) act as downstream effectors of the Hippo pathway, and their subcellular location and transcriptional activities are affected by multiple post- translational modifications (PTMs). After phosphorylation, YAP1/TAZ binds to the 14-3-3 protein, which induces cytoplasmic retention of YAP1/TAZ. Moreover, phosphorylated YAP1/TAZ is likely to undergo ubiquitination degradation that is dependent on the β -transducin repeat-containing E3 ubiquitin protein ligase complex. The phosphorylation of YAP at Ser127 promotes its cytoplasmic retention, whereas phosphorylation at Ser397 induces degradation. (PMID: 29682330, PMID: 26039999)		
Storage	Storage: Store at -80°C. The product is shipped with Storage Buffer: PBS Only	ice packs. Upon receipt, store it immediatel	y at -80°C

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Calyculin A treated and untreated HepG2 cells were subjected to SDS PAGE followed by western blot with 80694-2-RR (Phospho-YAP1 (Ser127) antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80694-2-PBS in a different storage buffer formulation. 1X10^6 HepG2 cells untreated (dashed lines) or treated with Calyculin A intracellularly stained with 0.13 ug Phospho-YAP1 (Ser127) Recombinant antibody (80694-2-RR, Clone:240779G4) and CoraLite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.13 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH. This data was developed