

For Research Use Only

c-MYC Recombinant antibody

Catalog Number: 80845-1-RR

Featured Product



Basic Information

Catalog Number:

80845-1-RR

Concentration:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG1263

GenBank Accession Number:

BC000141

GeneID (NCBI):

4609

UNIPROT ID:

P01106

Full Name:

v-myc myelocytomatosis viral
oncogene homolog (avian)

Calculated MW:

49 kDa

Observed MW:

50-65 kDa

Purification Method:

Protein A purification

CloneNo.:

2B17

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:200-1:800

IF/ICC 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : HEK-293 cells, A549 cells, HeLa cells, K-562 cells,
HL-60 cells

IHC : human colon cancer tissue,

IF/ICC : HeLa cells, HEK-293 cells, U2OS cells

Background Information

MYC contains one basic helix-loop-helix (bHLH) domain. This protein is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. It seems to activate the transcription of growth-related genes. It binds DNA in a non-specific manner, yet also specifically recognizes the core sequence 5'-CAC[GA]TG-3'. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 80845-1-RR (MYC antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MYC antibody (80845-1-RR, Clone: 2B17) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).




WB result of MYC antibody (80845-1-RR; 1:600; incubated at room temperature for 1.5 hours) with sh-Control and sh-MYC transfected A549 cells.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 80845-1-RR (c-MYC antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1×10^6 HeLa cells were intracellularly stained with 0.25 μ g c-MYC Recombinant antibody (80845-1-RR, Clone:2B17) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 μ g Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLI) kinetic assays of 80845-1-RR against Human c-MYC were performed. The affinity constant is 2.03 nM.