For Research Use Only

Osteopontin Recombinant antibody, PBS Only

Concentration: 1 mg/ml

Catalog Number: 80912-4-PBS



Purification Method:

Protein A purification

CloneNo.:

6B11

Basic Information

Catalog Number: 80912-4-PBS

BC007016 GeneID (NCBI): 6696

Source: UNIPROT ID:
Rabbit P10451-5
Isotype: Full Name:

lgG secreted phosphoprotein 1

Immunogen Catalog Number:Calculated MW:AG19216314 aa, 35 kDaObserved MW:

70 kDa, 44-66 kDa

GenBank Accession Number:

Applications

Tested Applications:

WB, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

Osteopontin (OPN), also known as SPP1, is a secreted glycophosphoprotein that belongs to the small integrinbinding ligand N-linked glycoprotein (SIBLING) family. Originally isolated from bone, OPN has been found in kidneys, vascular tissues, biological fluids, and various tumor tissues (PMID: 15138464; 16406521). OPN can interact with CD44 and integrins and regulate diverse biological processes. It has a multifaceted role in bone development and remodeling, and is also involved in the inflammatory and immune response, oncogenesis and cancer progression. The very acidic nature of OPN, as well as the presence of variable posttranslational modifications, has led to anomalous migration in SDS-polyacrylamide gels and therefore to reports of different molecular weights for OPN (PMID: 8293561). Depending on the cell and tissue source and/or the SDS-PAGE system, OPN migrates with a molecular weight of 44-80 kDa, as well as at some smaller bands corresponding to peptide fragments (PMID: 8195113; 17890765).

Storage

Storage:


Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 80912-4-RR (OPN, SPP1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80912-4-PBS in a different storage buffer formulation.



1x10^6 HepG2 cells were intracellularly stained with 0.25 ug OPN, SPP1 Recombinant antibody (80912-4-RR, Clone:6B11) and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 80912-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLL) kinetic assays of 80912-4-RR against Human OPN, SPP1 were performed. The affinity constant is 199 nM.