

For Research Use Only

# HIF-1 alpha Recombinant antibody, PBS Only

Catalog Number: 80933-1-PBS

Featured Product



## Basic Information

Catalog Number:

80933-1-PBS

Concentration:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG15198

GenBank Accession Number:

BC012527

GeneID (NCBI):

3091

UNIPROT ID:

Q16665

Full Name:

hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)

Calculated MW:

826 aa, 93 kDa

Observed MW:

120 kDa

Purification Method:

Protein A purification

CloneNo.:

2K1

## Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

human

## Background Information

HIF1α, the major regulator of the cellular responses to hypoxia, consists of an oxygen-sensitive subunit, HIF1α (HIF1A), and an oxygen-insensitive subunit, HIF1β (arylhydrocarbon receptor nuclear transporter [ARNT]). Under normal oxygen conditions, HIF1α is continuously produced and destroyed, in a process involving hydroxylation, interaction with von Hippel-Lindau (VHL) protein, polyubiquitylation and subsequent proteasomal degradation. Under hypoxic conditions, hydroxylation is impaired and HIF1α is stabilized. HIF1α localizes in cytoplasm in normoxia, but it can translocate into nuclear in response to hypoxia. The calculated molecular weight of HIF1α is 93 kDa, but the modified protein HIF1α is about 110-120kDa (PMID: 11698256, PMID: 7539918).

## Storage

Storage:

Store at -80°C.

**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

Storage Buffer:

PBS only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Untreated and Cobalt Chloride treated HeLa cells were subjected to SDS PAGE followed by western blot with 80933-1-RR (HIF 1α antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 80933-1-PBS in a different storage buffer formulation.




WB result of HIF-1 alpha antibody (80933-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HIF-1 alpha transfected HeLa cells. Sample 1: non-treated sh-Control transfected HeLa cells, Sample 2: Cobalt Chloride treated sh-Control transfected HeLa cells, Sample 3: Cobalt Chloride treated sh-HIF-1 alpha transfected HeLa cells. This data was developed using the same antibody clone with 80933-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed Cobalt Chloride treated HepG2 cells using HIF 1α antibody (80933-1-RR, Clone: 2K1 ) at dilution of 1:200 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was developed using the same antibody clone with 80933-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed Cobalt Chloride treated HeLa cells using HIF-1 alpha antibody (80933-1-RR, Clone: 2K1 ) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-phalloidin (red). This data was developed using the same antibody clone with 80933-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 80933-1-RR against Human HIF-1 alpha were performed. The affinity constant is 0.235 nM.