For Research Use Only

CD3 Recombinant antibody

Catalog Number:81324-1-RR

Featured Product



Basic Information

Catalog Number: GenBank Accession Number: 81324-1-RR BC049847 GeneID (NCBI):

Concentration: 500 ug/ml 916

ENSEMBL Gene ID: Source: Rabbit ENSG00000198851 UNIPROT ID: Isotype: P07766 Immunogen Catalog Number: Full Name:

AG11797

CD3e molecule, epsilon (CD3-TCR

Calculated MW: 207 aa, 23 kDa Observed MW: 23-25 kDa

complex)

Applications

Tested Applications: WB, IHC, IF-P, ELISA Species Specificity: human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: Jurkat cells, MOLT-4 cells, pig thymus tissue, rat thymus tissue, mouse thymus tissue

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:5000-1:20000

IF-P 1:50-1:500

CloneNo.:

4|19

IHC: human colon tissue, human tonsillitis tissue, mouse small intestine tissue, mouse spleen tissue

IF-P: mouse spleen tissue, mouse small intestine

Background Information

CD3 is a complex of proteins that directly associates with the T cell receptor (TCR). The TCR/CD3 complex of Tlymphocytes consists of either a TCR alpha/beta or TCR gamma/delta heterodimer coexpressed at the cell surface with the invariant subunits of CD3 labeled gamma, delta, epsilon, zeta, and eta. The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction. CD3 is considered to be a pan-T cell marker for detection of normal and neoplastic T cells. This antibody is directed against the epsilon chain of human CD3 molecule.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81324-1-RR (CD3E antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



WB result of CD3 antibody (81324-1-RR; 1:60000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CD3 transfected Jurkat cells.




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 81324-1-RR (CD3 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 81324-1-RR (CD3 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 81324-1-RR (CD3 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse spleen tissue using CD3 antibody (81324-1-RR, Clone: 4l19) at dilution of 1:200 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse small intestine tissue using CD3 antibody (81324-1-RR, Clone: 4l19) at dilution of 1:1000 and Coralite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 81324-1-RR against Human CD3 were performed. The affinity constant is 1.13 nM.