For Research Use Only

Phospho-TBC1D1 (Ser660) Recombinant antibody, PBS Only

Catalog Number:81363-2-PBS





Basic Information	Catalog Number: 81363-2-PBS	GenBank Accession Number: BC050321	Purification Method: Protein A purification
	Size: 1mg/ml	Genel D (NCBI): 23216	CloneNo.: 7N19
	Source: Rabbit	UNIPROT ID: Q86TIO	
	lsotype: IgG	Full Name: TBC1 (tre-2/USP6, BUB2, cdc16) domain family, member 1	
		Calculated MW: 133 kDa	
		Observed MW: 160 kDa	
Applications	Tested Applications: WB, Indirect ELISA		
	Species Specificity: human, mouse		
Background Information	TBC1D1 belong to the family of RabGAPs and its physiological roles in glucose homeostasis and its implication in human diseases have already been characterized in skeletal muscle and adipocytes in a large number of independent studies. TBC1D1 including twophospho-tyrosine-binding (PTB) domains, a calmodulin-binding domain(CBD) and the func-tional GAP domain responsible for catalyzing GTP hydrolysis. TBC1D1 is phosphorylated by AMPK at Ser660 and Thr237, and by AKT at Thr596. The RabGAPs TBC1D1 is important regulators of glucose homeostasis inskeletal muscle and adipose tissue. (PMID: 31627187, PMID: 18477703)		
Storage	Storage: Store at -80°C. The product is shipped with in Storage Buffer: PBS Only	ce packs. Upon receipt, store it immediatel	yat-80°C

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Non-treated and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 81363-2-RR (Phospho-TBC 1D1 (Ser660) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin antibody as loading control. This data was developed using the same antibody clone with 81363-2-PBS in a different storage buffer formulation.

Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 81363-2-RR (Phospho-TBC1D1 (Ser660) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin antibody as loading control. This data was developed using the same antibody clone with 81363-2-PBS in a different storage buffer formulation.