For Research Use Only

GLUT1 Recombinant antibody


Catalog Number:81463-1-RR 2 Publications




Basic Information	Catalog Number: 81463-1-RR	GenBank Accession Number: BC121804		Purification Method: Protein A purification	
	Size: 1000 ug/ml	GeneID (NC 6513	BI):	CloneNo.: 6F14	
	Source: Rabbit	UNIPROT ID P11166	t.	Recommended Dilutions: WB 1:5000-1:50000	
	Isotype: IgG Immunogen Catalog Number: AG16282	Full Name: solute carrier family 2 (facilitated glucose transporter), member 1		IHC 1:1000-1:4000	
			Calculated MW: 492 aa, 54 kDa		
		Observed M 45-55 kDa			
Applications	Tested Applications:	Positive Controls:			
	WB, IHC, ELISA Cited Applications:		WB : unboiled HEK-293 cells, HeLa cells (incubated 37°C)		
	WB, IHC Species Specificity:		IHC : human skin cancer tissue,		
	human Cited Species: human, mouse				
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	GLUT1, also known as SLC2A1, is a ubiquitously expressed glucose transporter and is responsible for the basal lever of glucose uptake in most cell types. Human erythrocytes express the highest level of GLUT1. Defects in SLC2A1 are the cause of GLUT1 deficiency syndrome type 1 and type 2. High expression of GLUT1 has been reported to be a reliable immunohistochemical marker for juvenile hemangiomas. GLUT1 protein may appear as two or more distinct forms among 43 kDa to 55 kDa due to the different glycosylation states. And the conversion of the highly glycosylated form of GLUT1 to a less glycosylated form has been reported to correlate with differentiation (PMID: 8263524, 23302780).				
Notable Publications	Author	Pubmed ID	Journal	Application	
	Zhao Cui	38622131	Cell Death Dis	WB,IHC	
	Jiajin Li	37311977	Br J Cancer	WB	

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded skin cancer slide using 81463-1-RR (GLUT1 antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Various lysates were subjected to SDS PAGE followed by western blot with 81463-1-RR (SLC2A1,GLUT1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.