

For Research Use Only

HDAC2 Recombinant monoclonal antibody, PBS Only

Catalog Number: 81497-4-PBS



Basic Information

Catalog Number: 81497-4-PBS	GenBank Accession Number: BC031055	Purification Method: Protein A purification
Source: Rabbit	GeneID (NCBI): 3066	CloneNo.: 252017B9
Isotype: IgG	UNIPROT ID: Q92769	
Immunogen Catalog Number: AG3607	Full Name: histone deacetylase 2	
	Calculated MW: 458 aa, 52 kDa; 488 aa, 55 kDa	
	Observed MW: 55-60 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, IP, Indirect ELISA

Species Specificity:
human, mouse

Background Information

Histone deacetylases (HDAC) are a class of enzymes that remove the acetyl groups from the lysine residues, leading to the formation of a condensed and transcriptionally silenced chromatin. Histone deacetylases act via the formation of large multiprotein complexes, and are responsible for the deacetylation of lysine residues at the N-terminal regions of core histones (H2A, H2B, H3, and H4). At least 4 classes of HDAC were identified. As a class, I HDAC, HDAC2 was primarily found in the nucleus. HDAC2 forms transcriptional repressor complexes by associating with many different proteins, including YY1, a mammalian zinc-finger transcription factor. Thus, it plays an important role in transcriptional regulation, cell cycle progression, and developmental events. This antibody is a rabbit monoclonal antibody raised against residues near the C terminus of human HDAC2.

Storage


Storage:
Store at -80°C.
The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:
PBS only, pH7.3


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81497-4-RR (HDAC2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 81497-4-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 81497-4-RR (HDAC2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 81497-4-PBS in a different storage buffer formulation.




IP result of anti-HDAC2 (IP:81497-4-RR, 4ug; Detection:81497-4-RR 1:800) with HEK-293 cells lysate 1080 ug. This data was developed using the same antibody clone with 81497-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HDAC2 antibody (81497-4-RR, Clone: 252017B9) at dilution of 1:1000 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 81497-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using HDAC2 antibody (81497-4-RR, Clone: 252017B9) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 81497-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HDAC2 antibody (81497-4-RR, Clone: 252017B9) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 81497-4-PBS in a different storage buffer formulation.