For Research Use Only

MMP2 Recombinant antibody

Catalog Number:81515-2-RR




Basic Information	Catalog Number: 81515-2-RR	GenBank Accession Number: BC002576	Purification Method: Protein A purfication
	Size: 300 ug/ml	GenelD (NCBI): 4313	CloneNo.: 240131F5
	Source: Rabbit	UNIPROT ID: P08253	Recommended Dilutions: IF/ICC 1:50-1:500
	Isotype: IgG Immunogen Catalog Number: AG0549	Full Name: matrix metallopeptidase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase) Calculated MW: 72 kDa	
		Applications	Tested Applications:
IF/ICC, FC (Intra), ELISA Species Specificity: human	IF/ICC : HepG2 cells,		
Background Information	MMP2, also named as CLG4A, Gelatinase Am, TBE-1 and PEX, belongs to the peptidase M10A family. It is ubiquitinous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. MMP2 contributes to myocardial oxidative stress by regulating the activity of GSK3beta. It cleaves GSK3beta in vitro. MMP2 can be cleaved into PEX chain(~60kd). Western blot analysis showed that the 72 kDa and 62/65 kDa proteinase activities were pro-MMP2 and the active enzyme, respectively (PMID:11112697,PMID: 30867536).		
Storage	Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage		


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using MMP2 antibody (81515-2-RR, Clone: 240131F5) at dilution of 1:100 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug MMP2 Recombinant antibody (81515-2-RR, Clone:240131F5) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 HT-29 cells were intracellularly stained with 0.25 ug MMP2 Recombinant Antibody (81515-2-RR, Clone:240131F5) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 81515-2-RR against Human MMP2 were performed. The affinity constant is below 1 pM.