For Research Use Only


## DRP1 (C-terminal) Recombinant antibody, PBS Only Catalog Number:81561-1-PBS Featured Product




Catalog Number: 81561-1-PBS	GenBank Accession Number: BC024590	Purification Method: Protein A purification	
Size: 1mg/ml	GenelD (NCBI): 10059	CloneNo.: 2A19	
Source: Rabbit	UNIPROT ID: O00429		
Isotype: IgG	Full Name: dynamin 1-like		
Immunogen Catalog Number: AG3644	Calculated MW: 736 aa, 82 kDa		
	Observed MW: 70-80 kDa		
Tested Applications: WB, IHC, IF/ICC, FC (Intra), Indired	Tested Applications: WB, IHC, IF/ICC, FC (Intra), Indirect ELISA		
Species Specificity: human, mouse, rat			
of large GTPases that mediate me important role in the division of g membrane fission in mammalian	DNM1L(Dynamin-1-like protein), also known as dynamin-related protein 1 (DRP1), belongs to the dynamin family of large GTPases that mediate membrane remodeling during a variety of cellular processes. DNM1L has an important role in the division of growing mitochondria and peroxisomes and also mediates outer mitochondrial membrane fission in mammalian cells. DNM1L is ubiquitously expressed with abundant expression in skeletal muscle, heart, kidney and brain.Variable isoforms of DNM1L have been reported in a tissue-specific manner due to the alternative splicing.		
Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediatel	yat-80°C	
	81561-1-PBS Size: Img/ml Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG3644 Tested Applications: WB, IHC, IF/ICC, FC (Intra), Indired Species Specificity: human, mouse, rat DNM1L(Dynamin-1-like protein), of large GTPases that mediate me important role in the division of g membrane fission in mammalian muscle, heart, kidney and brain.V. the alternative splicing. Storage: Storage: Storage Storage Buffer:	81561-1-PBS BC024590   Size: GeneID (NCBI):   1mg/ml 10059   Source: UNIPROT ID:   Rabbit O00429   Isotype: Full Name:   IgG dynamin 1-like   Immunogen Catalog Number: Calculated MW:   AG3644 736 aa, 82 kDa   Observed MW: 70-80 kDa   Tested Applications: WB, IHC, IF/ICC, FC (Intra), Indirect ELISA   Species Specificity: human, mouse, rat   ON DNM11(Dynamin-1-like protein), also known as dynamin-related protect of large GTPases that mediate membrane remodeling during a variety of important role in the division of growing mitochondria and peroxisome membrane fission in marmalian cells. DIM1L is ubiquitously express muscle, heart, kidney and brain.Variable isoforms of DNM1L have been the alternative splicing.   Storage: Storage Listing.   Storage Buffer: Storage Buffer:	

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 81561-1-RR (DRP1 (C-terminal) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 81561-1-PBS in a different storage buffer formulation.



WB result of DRP1 (C-terminal) antibody (81561-1-RR; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-DRP1 (C-terminal) transfected HEK-293 cells. This data was developed using the same antibody clone with 81561-1-PBS in a different storage buffer formulation.




1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human DRP1 (C-terminal) (81561-1-RR, Clone:2A19) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 81561-1-PBS in a different storage buffer formulation.






Various lysates were subjected to SDS PAGE Immunohistochemical analysis of paraffinfollowed by western blot with 81561-1-RR (DRP1 (C-terminal) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This embedded mouse cerebellum tissue slide using 81561-1-RR (DRP1 (C-terminal) antibody) at dilution of 1:1000 (under 10x lens). Heat mediated data was developed using the same antibody clone with 81561-1-PBS in a different storage buffer antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 81561-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 81561-1-RR (DRP1 (C-terminal) antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 81561-1-PBS in a different storage buffer formulation.



formulation.

Immunofluorescent analysis of (-20°C Methanol) fixed NIH/3T3 cells using DRP1 (C-terminal) antibody (81561-1-RR, Clone: 2A19) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L). This data was dwolened using the came antibody clone with developed using the same antibody clone with 81561-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 81561-1-RR against Human DRP1 (C-terminal) were performed. The affinity constant is 0.825 nM.

