## For Research Use Only

## SNAI1 Recombinant antibody

Catalog Number:81584-5-RR



**Purification Method:** 

Protein A purfication

WB 1:5000-1:50000

Recommended Dilutions:

CloneNo.:

240164C11

**Basic Information** 

Catalog Number: GenBank Accession Number:

81584-5-RR BC012910 GeneID (NCBI): Size: 1000 ug/ml 6615 **UNIPROT ID:** Source: Rabbit 095863

Full Name: Isotype:

snail homolog 1 (Drosophila)

Calculated MW: Immunogen Catalog Number: AG24248 264 aa, 29 kDa Observed MW:

30 kDa

**Applications** 

**Tested Applications:** WB, FC (Intra), ELISA Species Specificity:

human

Positive Controls:

WB: A549 cells, MCF-7 cells, BxPC-3 cells

## **Background Information**

SNAI1, a member of SNAI1 family of protein, participates in the epithelial to mesenchymal transition(EMT) and formation and maintenance of embryonic mesoderm. The snail family share a common structural, that a highly conserved C-terminal region containing a zinc finger transcription factor. SNAI1 interacts with other corepressor, such as Ajuba, PRMT5 and SIN3a or HDAC1 and 2, to repress the target gene. As the phosphorylation modification of SNAI1 protein, the range of molecular weight of SNAI1 is about 25-30 kDa (PMID: 22276203). Once phosphorylated (probably on Ser-107, Ser-111, Ser-115 and Ser-119) it is exported from the nucleus to the cytoplasm where subsequent phosphorylation of the destruction motif and ubiquitination involving BTRC occurs.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



Various lysates were subjected to SDS PAGE followed by western blot with 81584-5-RR (SNAI1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1x10^6 A549 cells were intracellularly stained with 0.25 ug SNAl1 Recombinant antibody (81584-5-RR, Clone:240164C11) and APC-Conjugated AffiniPure Goat Anti-Rabbit I gG(H+L)(red), or 0.25 ug I sotype Control (blue). Cells were fixed and permeabilized with True-Nuclear Transcription Factor Buffer Set.



Biolayer interferometry (BL1) kinetic assays of 81584-5-RR against Human SNAI1 were performed. The affinity constant is 0.64 nM.