For Research Use Only

Phospho-p70(S6K) (Thr229) Recombinant antibody




Catalog Number:81592-1-RR

Basic Information	Catalog Number: 81592-1-RR	GenBank Accession Number: BC053365	Purification Method: Protein A purification	
	Size: 1000 µg/ml	GenelD (NCBI): 6198	CloneNo.: 2G19	
	Source: Rabbit	UNIPROT ID: P23443	Recommended Dilutions: WB 1:5000-1:50000	
	Isotype: IgG	Full Name: ribosomal protein S6 kinase, 70kDa, polypeptide 1 Calculated MW: 59 kDa Observed MW: 65-85 kDa		
Applications	Tested Applications:	Positive Controls:		
	WB, ELISA WB : IGF-1 treated MCF-7 cells, Species Specificity: Human			
Background Information	RPS6KB1(Ribosomal protein S6 kinase beta-1) is also named as STK14A, p70 S6KA, and belongs to the S6 kinase subfamily. RPS6KB1 is a major substrate of mTOR and acts as a crucial effector of mTOR signaling pathway. It plays a key role in cell growth and proliferation by regulating INS sensitivity, metabolism, protein synthesis, and cell cycle. RPS6KB1 may play an important role in the progression of HCC and could serve as a potential molecular target for HCC therapy (PMID:22684641). The Rps6kb1 gene encodes the 70 kDa ribosomal protein S6 kinase (p70S6K). The PI3K/mTOR signalling pathway is one of the major mechanisms for controlling cell survival, proliferation, and metabolism and is the central regulator of translation of some components of the protein synthesis system. Due to alternative translation, two isoform S6K1 proteins are known to exist in mammalian cells: p85 S6K1 and p70 S6K1, which is identical to p85 S6K but lacks its first 23 amino acids. In addition, mammalian cells express a second S6K1 isoform spanning 316 amino acids (p31 S6K1).			
Storage	Storage Buffer:	0.02% sodium azide and 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated and IGF-1 treated MCF-7 cells were subjected to SDS PAGE followed by western blot with 81592-1-RR (Phosoho-p70(S6K) (Thr229) antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with 60004-1-Ig (GAPDH antibody) as loading control.