## For Research Use Only

## JNK Recombinant antibody

Catalog Number:81629-1-RR



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC130572

**Purification Method:** Protein A purification

81629-1-RR Size:

GeneID (NCBI):

CloneNo.:

1000 µg/ml

**UNIPROT ID:** 

6]16

Source: Rabbit

P45983 Full Name: Recommended Dilutions: WB 1:20000-1:100000

Isotype:

mitogen-activated protein kinase 8

IHC 1:50-1:500 IF/ICC 1:200-1:800

Immunogen Catalog Number: AG1639

Calculated MW: 48 kDa

Observed MW:

44-48 kDa, 50-55 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), ELISA

Positive Controls:

Species Specificity:

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, NIH/3T3 cells, HSC-T6 cells

human, mouse, rat

IHC: human colon cancer tissue, IF/ICC: HeLa cells, Jurkat cells

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

## **Background Information**

MAPK8(Mitogen-activated protein kinase 8) is also named as JNK1, PRKM8, SAPK1, SAPK1C and belongs to the MAP kinase subfamily. MAPK8 is activated by dual phosphorylation at a Thr-Pro-Tyr motif during response to UV light. MAPK8 functions to phosphorylate c-Jun at N-terminal serine regulatory sites of Ser-63 and Ser-73, mapping within the transactivation domain. Phosphorylation of these sites in response to UV results in transcriptional activation of c-Jun. It has some isoforms produced by alternative splicing with the molecular weight of 46 kDa and 48 kDa. This protein can be phosphorylated and this antibody recognizes the 46 kDa and 55 kDa bands in western blot(PMID:11062067). This antibody can recognize JNK1, JNK2 and JNK3.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 81629-1-RR (JNK antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 81629-1-RR (JNK antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 81629-1-RR (JNK antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed Hela cells using JNK antibody (81629-1-RR, Clone: 6J16) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using JNK antibody (81629-1-RR, Clone: 6J16) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human JNK (81629-1-RR, Clone:6J16) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit I gG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLL) kinetic assays of 81629-1-RR against Human JNK were performed. The affinity constant is below 1 pM.