For Research Use Only

STAT1 Recombinant antibody, PBS Only

Catalog Number:82016-1-PBS

Featured Product




Basic Information	Catalog Number: 82016-1-PBS	GenBank Accession Number: BC002704	Purification Method: Protein A purification
	Size: 1mg/ml	GenelD (NCBI): 6772	CloneNo.: 2F16
	Source: Rabbit	UNIPROT ID: P42224	
	Isotype: IgG Immunogen Catalog Number: AG0199	Full Name: signal transducer and activator of transcription 1, 91kDa Calculated MW: 83 kDa	
		Applications	Tested Applications: WB, IHC, IF/ICC, IP, Indirect ELISA
Species Specificity: human, mouse, rat			
Background Information	STAT1 (signal transducers and activators of transcription 1) is a member of the STAT protein family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. Various ligands, including interferons- and -, EGF, PDGF and IL-6, can activate STAT1. STAT1 protein mediates the expression of a variety of genes, which is thought to be important for cell viability in response to different cell stimuli and pathogens.		
Storage	Storage: Store at -80°C. The product is shipped with ice pac Storage Buffer: PBS Only	:ks. Upon receipt, store it immediatel	y at -80℃


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82016-1-RR (STAT1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82016-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 82016-1-RR (STAT1 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82016-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 82016-1-RR (STAT1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82016-1-PBS in a different storage buffer formulation.




WB result of STAT1 antibody (82016-1-RR; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-STAT1 transfected HEK-293 cells. This data was developed using the same antibody clone with 82016-1-PBS in a different storage buffer formulation.



IP result of anti-STAT1 (IP:82016-1-RR, 4ug; Detection:82016-1-RR 1:10000) with PC-3 cells lysate 2000 ug. This data was developed using the same antibody clone with 82016-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed IFN alpha treated HeLa cells using STAT1 antibody (82016-1-RR, Clone: 2F16) at dilution of 1:1000 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-phalloidin (red). This data was developed using the same antibody clone with 82016-1-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 82016-1-RR against Human STAT1 were performed. The affinity constant is 0.501 nM.